

Specimen Label

Akari® 5SC Miticide/Insecticide



Suspension Concentrate

FOR USE ON GREENHOUSE AND NURSERY
ORNAMENTALS, NON-BEARING FRUIT TREES AND
VINES, CHRISTMAS TREES, AND INTERIOR
ORNAMENTAL PLANTINGS AND PLANTSCAPES

Active Ingredient

<i>Tert</i> -butyl(E)- α -(1,3-dimethyl-5-phenoxy)pyrazol -4-ylmethylenaminoxy)-p-toluate	5.0%
Other Ingredients.....	95.0%
TOTAL.....	100.0%

Contains 0.42 lb. active ingredient per U.S. gallon.

Keep Out of Reach of Children WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles or face shield). Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Harmful if inhaled. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment (PPE) Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Protective eyewear
- Waterproof gloves
- Shoes plus socks

First Aid

If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.

EMERGENCY NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC at 1-800-535-5053**.

Note to Physician: There is no specific antidote. All treatment should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

Refer to label booklet for additional precautionary information and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer," "Inherent Risks of Use," and "Limitation of Remedies" inside label booklet.**

For additional information on our products, please visit www.sepro.com.

EPA Reg. No. 71711-4-67690
EPA Est. No. 37429-GA-2
FPL 012406

Akari is a registered trademark of Nichino America, Inc.
Manufactured for: **SePRO Corporation** Carmel, IN 46032, U.S.A.

User Safety Recommendations

Do not eat, drink, smoke, or chew gum or tobacco while handling this product and until hands and face are thoroughly washed with soap and water. Do not use the toilet before thoroughly washing hands. If this product penetrated through your clothing or personal protective equipment, stop handling this product immediately, remove the clothing and equipment, wash your body thoroughly, and put on clean clothing and equipment before resuming the handling activity. After handling this product, remove personal protective equipment immediately. Wash the outside of gloves before taking them off. Shower or wash thoroughly and change into clean clothing as soon as possible. Discard clothing and personal protective equipment that cannot be reused, including clothing and other absorbent materials that have been drenched or thoroughly contaminated with this product's concentrate. Otherwise, wash clothing and personal protective equipment (including both the inside and outside of gloves) before each day of reuse according to manufacturers' directions or, if no such directions, in detergent and hot water. Keep and wash them separately from other laundry.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to fish and aquatic invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or Tribe, consult the agency in your State responsible for pesticide regulation.

GENERAL INFORMATION

Akari® 5SC is a contact miticide used for the control of spider mites and other mite pests of ornamentals. This product should be used in a program with other products to provide season-long protection. Rotate to one or more miticides utilizing a different mode of action (MOA) before reapplying Akari 5SC. The mode of action of Akari 5SC is a Mitochondrial Electron Transport Inhibitor (METI). Apply as a spray as set forth in the *Application Directions* section of this label, using sufficient water volume to obtain thorough coverage of plants. Under severe mite conditions, use the maximum rates and the shorter spray interval as specified on the label. Dense foliage or excessive growth will often prevent adequate coverage; adjust spray volumes accordingly.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Protective eyewear
- Waterproof gloves
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, or greenhouses. For other uses, including interiorscapes and other nonagricultural uses, do not enter treated areas without protective clothing until sprays have dried.

Akari 5SC is a 5% Suspension Concentrate containing 0.42 lb. of fenpyroximate per gallon, which, when mixed with water according to the *Directions for Use*, will control mites on ornamentals.

Akari 5SC works primarily through contact action, so good spray coverage is necessary. Mix with sufficient water and apply as a foliar spray to obtain uniform coverage. Treat plants when pests are immature or at a susceptible stage and populations are building, before crop damage occurs.

Akari 5SC stops mite damage immediately after application. Akari 5SC provides this stop-feeding action together with a cessation of egg laying, and mites die in 4 to 7 days. When Akari 5SC is used to control mites, suppression of mealybugs on the treated crop may be achieved. This product must be used in a program with other products to provide protection against mealybugs.

COMPATIBILITY OF SPRAY MIXTURES

Akari 5SC is believed to be compatible with most commonly used fungicides, insecticides, micronutrients, growth regulators, and spray adjuvants. Consult specific product labels for additional information. It is always a good idea to conduct a tank-mix compatibility test when you plan to mix Akari 5SC with other products.

To determine the physical compatibility of Akari 5SC with other products, use a jar test. Using a jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add dry formulations first, then flowables, then emulsifiable concentrates last. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank. It is suggested that combinations be used on a small number of plants before treating large areas to check for cultivar sensitivity.

MIXING DIRECTIONS

Shake well before using

Please read *Compatibility of Spray Mixtures* section prior to any tank mixing with Akari 5SC. Begin with clean equipment. Prepare only the amount of spray solution needed to treat the target area. Add sufficient clean water to the spray tank for one-half of the mix load. Where possible, start agitation and properly suspend the necessary amount of Akari 5SC in the tank. Agitate to ensure thorough mixing while adding the remaining required water volume and other products. Akari 5SC should be properly suspended and diluted prior to the addition of any adjuvant or before spraying. Consult the adjuvant label or manufacturer for crop tolerance and safety information when using Akari 5SC. Maintain agitation during mixing and application. If agitation is stopped for any reason, the spray solution must be thoroughly re-mixed prior to further use.

APPLICATION DIRECTIONS

Akari 5SC is recommended for use on all ornamental crops including flowering and foliage crops, nursery crops, non-bearing fruit trees and vines, and Christmas trees. These include those crops grown in greenhouses, shade and lathe houses, outdoor nurseries and fields, and in interiorscapes. The use directions of this product are based on the results of product testing programs on a wide variety of ornamental plants. However, it is impossible to test this product on all species and cultivars. The phytotoxicity of Akari 5SC has been assessed on a wide variety of common ornamental plants with no phytotoxicity. However, not all plant species and their varieties and cultivars have been tested with possible tank mix combinations, sequential pesticide treatments, and adjuvants and surfactants. Local conditions also can influence crop tolerance and may not match those under which testing has been conducted. Therefore, before using Akari 5SC, test the product on a sample of the crop to be treated to ensure that a phytotoxic response will not occur as a result of applications. Applications should be made immediately after the spray solution is prepared. Thorough spray coverage is essential for mite control.

Applications may be made with high or low volume spray equipment that provides thorough coverage of the plant. Apply with properly calibrated spray equipment. A wetting agent or other spray adjuvant, approved for use on the crop, may be added to spray solutions according to the manufacturer's use instructions to achieve optimum mite control.

Table 1: Recommended Application Rates for Akari 5SC on Ornamental Trees, Vines, Shrubs, Foliage and Flowering Plants

Pest	Application Rate Fl. Oz./100 Gallons	Specific Directions
Spider mites	16 - 24 ounces	Use the lower rate for low to moderate mite populations.
Tarsonemid mites • Broad mite (<i>Polyphagotarsonemus latus</i>) • Cyclamen mite (<i>Phytonemus pallidus</i>)	24 ounces	Use in sufficient volume to obtain uniform plant coverage. 100 gallons of spray will typically cover 20,000 sq. ft. of greenhouse.
Eriophyid mites (rust, gall, and bud mites) including but not limited to: • Hemlock rust mite (<i>Nalepella tsugifoliae</i>) • Hackberry rust mite	24 ounces	Do not apply more than 10 gallons of spray per 1000 sq. ft. per application. Make sequential application of a miticide, with a different mode of action, up to 21 days after initial Akari 5SC application.
Mealybugs, including but not limited to: • Citrus mealybug (<i>Planococcus citri</i>) • Long-tailed mealybug (<i>Pseudococcus longispinus</i>)	24 ounces	For suppression of certain mealybug pests, use in sufficient volume to obtain uniform plant coverage. 100 gallons of spray will typically cover 20,000 sq. ft. of greenhouse. Do not apply more than 10 gallons of spray per 1000 sq. ft. per application.

Akari 5SC is recommended for use on all ornamental crops and plantings including those listed in Table 2.

Table 2: List of Plant Species Tested for Tolerance to Akari 5SC¹

Common Name	Scientific Name
African violet	<i>Saintpaulia ionantha</i>
Azalea	<i>Rhododendron</i> spp.
Chrysanthemum	<i>Chrysanthemum morifolium</i>
Citrus, nonbearing	<i>Citrus</i> spp.
Coleus	<i>Coleus X hybridus</i>
Cotoneaster	<i>Cotoneaster</i> spp.
Dieffenbachia	<i>Dieffenbachia</i> spp.
English ivy	<i>Hedera helix</i>
Fern	<i>Davallia</i> spp.
Ficus	<i>Ficus</i> spp.
Geranium	<i>Geranium</i> spp.
Gerbera	<i>Gerbera</i> spp.
Grape, nonbearing	<i>Vitis</i> spp.
Hackberry	<i>Celtis occidentalis</i>
Hemlock	<i>Tsuga</i> spp.
Holly (e.g. blue, Burford, Japanese)	<i>Ilex</i> spp.
Hypoestes	<i>Hypoestes</i> spp.
Juniper (e.g. creeping, Parsons, upright)	<i>Juniperus</i> spp.
Marigold	<i>Tagetes erecta</i>
New Guinea impatiens	<i>Impatiens linearifolia</i>
Pittosporum	<i>Pittosporum</i> spp.
Poinsettia	<i>Euphorbia pulcherrima</i>
Pyracantha (firethorn, red firethorn)	<i>Pyracantha angustifolia</i> , <i>P. coccinea</i>
Rose, cut and miniature	<i>Rosa</i> spp.
Schefflera	<i>Schefflera</i> spp.
Sweetpotato vine, ornamental	<i>Ipomoea batatas</i>
Zinnia	<i>Zinnia elegans</i>

¹Local conditions can influence crop tolerance and may not match those under which these species were tested. Before using Akari 5SC, test the product on a small sample of the crop to be treated.

GENERAL USE LIMITATIONS AND PRECAUTIONS

- In Florida, do not use on bearing or nonbearing commercial fruit trees and vines.
- **Do not** apply more than 48 fl. oz. per crop cycle or per growing season, whichever is longer.
- **Do not** apply this product through any type of irrigation system.
- **Do not** apply this product as a smoke or aerosol.
- **Do not** apply this product to plants used for food or animal feed.
- **Do not** apply to fruit trees and vines that will bear harvestable fruit within 12 months.
- **Do not** enter the entire enclosed area (treated area) without protective equipment for 12 hours unless one of the following items is completed:
 1. Ten air exchanges
 2. Two hours of system ventilation
 3. Four hours of ventilation using vents, windows, or other passive ventilation
 4. Eleven hours with no ventilation followed by 1 hour of mechanical ventilation
 5. Eleven hours with no ventilation followed by 2 hours of passive ventilation
 6. All required PPE is worn.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse, or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Resistance Management

DO NOT use Akari 5SC in successive miticide applications in the same production area. Allow at least 2 different chemicals with different modes of action to be used between treatments of Akari 5SC. Consult your local Horticultural Advisor for the most appropriate alternative products. Resistance management strategies should include alternation or tank mixtures with a miticide currently recommended for mite control. Resistance management strategies recommend that you **DO NOT** apply rates lower than recommended on the label. For best results, make no more than two applications of Akari 5SC per growing season, or per crop cycle, whichever is longer.

IMPORTANT: READ WARRANTY STATEMENT BEFORE USE

Warranty Disclaimer

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc. abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation as the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" can not be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or "Limitations of Remedies" in any manner.