

Specimen Label

Talus[®] 40SC Insect Growth Regulator



For Greenhouse and Outdoor Ornamentals and Greenhouse Tomatoes

Active Ingredient
Buprofezin:
2-tert-butylimino-3-isopropyl-5-phenylperhydro-
1,3,5-thiadiazin-4-one 40.00%
Other Ingredients 60.00%
TOTAL 100.00%

Contains 3.6 lbs. buprofezin per U.S. gallon.

Keep Out of Reach of Children WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes substantial but temporary eye injury. Avoid contact with skin or clothing. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant (such as nitrile or butyl) gloves
- Protective eyewear
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- Remove and wash contaminated clothing before reuse.

First Aid	
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call poison control center or doctor for treatment advice.
If on skin	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.

EMERGENCY NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC at 1-800-535-5053**.

Note to Physician: There is no specific antidote. All treatment should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

Refer to inside of label booklet for additional **Precautionary Information** and **Directions for Use** including **Storage and Disposal**.

Notice: Read the entire label before using. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer", "Inherent Risks of Use" and "Limitation of Remedies" inside label booklet.

For additional information on our products, please visit www.sepro.com.

EPA Reg. No. 71711-20-67690
FPL 061206

*Talus is a trademark of Nichino America, Inc.
Manufactured for: **SePRO Corporation** Carmel, IN 46032 U.S.A.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4–6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant (such as nitrile or butyl) gloves
- Protective eyewear
- Shoes plus socks.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, or greenhouses. For other uses, including golf courses and other nonagricultural uses, do not enter treated areas without protective clothing until sprays have dried.

GENERAL INFORMATION

Talus 40SC insect growth regulator contains the active ingredient, buprofezin. It is effective against the nymph stages of whitefly, scales, mealybugs, planthoppers, and leafhoppers by inhibiting chitin biosynthesis, suppressing oviposition of adults, and reducing viability of eggs. Treated susceptible pests may remain alive on the plant for 3 - 7 days, but feeding damage during this time is typically very low. Talus 40SC is not disruptive to beneficial insects and mites.

Good coverage is essential. Apply by ground or air in sufficient water volume. Orient nozzles to assure good coverage. Use of higher volume of water will assure better coverage, especially under adverse conditions such as hot, dry weather, and/or a dense canopy. The entire field should be treated.

Apply when economic infestations occur based on local information.

Talus 40SC is not for sale, sale into, distribution, and or use in Nassau and Suffolk counties of New York State.

INSECTS CONTROLLED

Whiteflies: Silverleaf whitefly; Sweet potato whitefly; Greenhouse whitefly; Ash whitefly

Mealybugs: Grape mealybug; Vine mealybug; Citrus mealybug; Comstock mealybug; Longtailed mealybug; Mexican mealybug; Obscure mealybug

Leafhoppers and Planthoppers: Eastern grape leafhopper; Glassy-winged sharpshooter; Variegated leafhopper; Western grape leafhopper; Cherry leafhopper; Potato leafhopper; Brown planthopper

Scales:

Soft Scales: Citricola scale; Barnacle scale; Black scale; Brown soft scale; False oleander scale; Hemispherical scale; Indian wax scale and other wax scales; Tessellated scale; White peach scale

Armored Scales: Boisduval scale; Cactus scale; California red scale; Coconut scale; Fern scale; Florida red scale; Oystershell scale; San Jose scale

Margarodid scale: Cottony cushion scale

RESISTANCE MANAGEMENT

Talus 40SC belongs to a class of chemicals not previously used as pesticides and not known to be cross-resistant to other classes of insecticides. However, insect pests are known to develop resistance to products used repeatedly. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotating with insecticides with different modes of action. Consult your local pest control advisor or extension office for details. If resistance to this

product develops in your area, it may not provide adequate control. If you experience difficulty with control, and resistance is a likely cause, consult your local state university horticultural specialist or local agricultural authorities for the best alternative method of control. To preserve the usefulness of Talus 40SC, do not make more than two consecutive applications. Prior to subsequent applications, use an alternative chemistry with a different mode of action. If a different crop in an adjacent field has been treated with Talus 40SC for whitefly control within the past 28 days, do not apply again as the first whitefly application. Always consult your local crop advisor for the most appropriate control decision for your area. Resistance management strategies advise against applying rates lower than those recommended on the label.

For greenhouses: If another insect growth regulator with the mode of action as a chitin synthesis inhibitor (i.e. cyromazine, diflubenzuron, and novaluron) has been used in the greenhouse within 28 days, do not apply Talus 40SC as the first whitefly application.

USE PRECAUTIONS

- Apply in a minimum finished spray volume of five (5) gallons per acre by air or twenty (20) gallons per acre by ground, unless otherwise specified in Directions for Use.
- **Crop Rotational Restrictions:** All crops listed on this label may be planted at any time. Small grain crops and leafy vegetables may be planted 30 days following application. All other crops may be planted 60 days following application.
- **For greenhouses:** Apply in a minimum of twenty (20) gallons per acre. Be sure to apply in sufficient water to obtain complete coverage of all plant parts. Applications may be made with high volume, low volume, or ultra low volume (thermal and non-thermal foggers, misters, etc.) ground equipment. Follow the spray equipment manufacturer's directions to determine the amount of spray solution required to obtain thorough coverage. Consult the spray equipment manufacturer's operator's manual, spray nozzle catalogs and/or your crop advisor for more information.
- **Do not apply this product through any type of irrigation system.**

Application Rate Chart for Talus 40SC Insect Growth Regulator

Crop	Pest	Formulated Product/A	Recommendations and Use Restrictions
Greenhouse Tomatoes	Leafhoppers Mealybugs Whiteflies	13.6 fl oz 13.6 fl oz/A 9 to 13.6 fl oz	<ul style="list-style-type: none"> • Treatment should be applied when population level reaches economic threshold. Consult local and state agricultural authorities for details. • For greenhouse tomatoes, apply by ground on 2 acres as a minimum with 20 gallons water per acre. Apply in sufficient water to obtain complete coverage of all plant parts. Applications may be made with high volume, low volume or ultra low volume (thermal and non-thermal foggers, misters, etc.) ground equipment only. • Follow the spray equipment manufacturer's directions to determine the amount of spray solution required to obtain thorough coverage. Consult the spray equipment manufacturer's operator's manual, spray nozzle catalogs and/or your crop advisor for more information. • Apply no more than two applications per season. • Allow at least 28 days between applications. • Do not apply within 7 days of harvest.
Ornamental plants in greenhouses; lath and shadehouses; nurseries; landscape ornamentals; ground covers; field- and container-grown ornamentals; non-bearing fruit and nut trees in nurseries; Christmas trees	Leafhoppers Mealybugs Scales Whiteflies	18 fl oz/100 gal 18 fl oz/100 gal 21.5 fl oz/100 gal 9 fl oz/100 gal	<ul style="list-style-type: none"> • Apply the specified dosage as a foliar spray in sufficient water for complete, uniform coverage, including stems and underside of leaves. Spray to the point of runoff. • For whitefly or leafhopper control: Make first application as soon as adult insects begin to appear. • For mealybug control: Make first application as soon as insect activity is observed. • For scale control: Make first application when crawlers are emerging. • If additional insecticide applications are required for control, use another class of chemistry or a different Insect Growth Regulator (IGR) with a different mode of action before making a second application of Talus 40SC. • Make no more than two applications of Talus 40SC per year. • Consult local or state agricultural authorities for details concerning economic thresholds for each target pest.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Store in original container, unopened, in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: DO NOT reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

Warranty Disclaimer

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SePRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks Of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner or application, or other factors, all of which are beyond the control of SePRO Corporation as the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies can not be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitations of Remedies in any manner.